AMENDMENTS TO THE CLAIMS

- 1. (currently amended) An isolated nucleic acid sequence which encodes a polypeptide and which is composed of a combination of the nucleic acid sequences of a biosynthetic nucleic acid sequence encoding of the a fatty acid or lipid metabolism enzyme and one of the following nucleic acids:
 - 1. a nucleic acid sequence with the sequence shown in SEQ ID NO: 1,
 - nucleic acid sequences which are derived from the nucleic acid sequence shown in SEQ ID NO: 1 as the result of the degeneracy of the genetic code,
 - 3. derivatives of the nucleic acid sequence shown in SEQ ID NO: 1 which encode polypeptides with the amino acid sequences shown in SEQ ID NO: 2 and which have at least 60% homology at the amino acid level,
 - 4. a nucleic acid sequence with the sequence shown in SEQ ID NO: 3 or the aminoterminal portion of the coding region of the sequence.
- 2. (currently amended) An isolated nucleic acid sequence as claimed in claim 1, wherein a sequence of the following protein groups is used as biosynthesis gene nucleic acid sequence of the fatty acid or lipid metabolism[[.]]:
 Acyl-CoA dehydrogenase(s), Acyl-ACP[=acyl carrier protein] desaturase(s), Acyl-ACP

thioesterase(s), fatty acid acyltransferase(s), fatty acid synthase(s), fatty acid hydroxylase(s), acetyl-coenzyme A carboxylase(s), acyl-coenzyme A oxidase(s), fatty acid desaturase(s), fatty acid acetylenases, lipoxygenases, triacylglycerol lipases,

allenoxide synthases, hydroperoxide lyases and/or fatty acid elongase(s).

- 3. (currently amended) An isolated nucleic acid sequence as claimed in claim1, wherein a sequence of the following protein groups is used as biosynthesis gene nucleic acid sequence of the fatty acid or lipid metabolism:
 fatty acid acryltransferase(s), Δ4 desaturase, Δ5 desaturase, Δ6 desaturase, Δ9 desaturase,
 Δ12 desaturase, Δ15 desaturase or a fatty acid elongase.
- 4. (currently amended) An isolated nucleic acid sequence as claimed in claim 1, wherein the derivatives mentioned under (c) have a homology at the amino acid level of 70%, preferably 80%[[, especially preferably of 90%]], over the entire region of the sequence shown in SEQ ID NO: 2 (Program PileIp, J. Mol. Evolution., 25, 351-360, 1987, Higgins et al., CABIOS, 5 1989: 151-153).
- 5. (original) An amino acid sequence encoded by a nucleic acid sequence as claimed in claim 1.
- 6. (currently amended) An isolated nucleic acid construct comprising a nucleic acid sequence as claimed in claim 1, wherein the nucleic acid sequence is linked to one or more regulatory signals.

- 7. (previously presented) The use if a nucleic acid sequence as claimed in claim 1 or of a nucleic acid construct comprising said nucleic acid sequence wherein the nucleic acid sequence is linked to one or more regulatory signals for the generation of transgenic plants.
- (previously presented) A vector comprising a nucleic acid sequence as claimed in claim
 1 or a nucleic acid construct comprising said nucleic acid sequence wherein the nucleic
 acid sequence is linked to one or more regulatory signals.
- 9. (original) A vector as claimed in claim 8, which is a linear or circular DNA, a phage, a virus, a transposon, a IS element, a phasmid, a phagemid, a cosmid or a plasmid.
- 10. (currently amended) An organism comprising at least one <u>isolated</u> nucleic acid sequence as claimed in claim 1, at least one <u>isolated</u> nucleic acid construct comprising said nucleic acid sequence wherein the nucleic acid sequence is linked to one or more regulatory signals or at least one vector comprising said nucleic acid sequence wherein the nucleic acid sequence is linked to one or more regulatory signals or a nucleic acid construct comprising said nucleic acid sequence wherein the nucleic acid sequence is linked to one or more regulatory signals.

- 11. (original) An organism as claimed in claim 10, which is a eukaryotic organism.
- 12. (previously presented) An organism as claimed in claim 10, which is a plant, a eukaryotic microorganism or an animal.
- (previously presented) An organism as claimed in claim 10, which is a plant, a fungus or a yeast.
- 14. (currently amended) An organism as claimed in claim 10, which is Yarrowie Iypolytica,

 Saccharomyces cereviseae, Traustochytrium, Arabidopsis thaliana, Brassica napus or

 Linium usitatissimum Yarrowie lypolytica, Saccharomyces cereviseae, Traustochytrium,

 Arabidopsis thaliana, Brassica napus or Linium usitatissimum.
- 15. (previously presented) A transgenic plant comprising a nucleic acid sequence as claimed in claim 1 or a nucleic acid construct comprising said nucleic acid sequence.

- 16. (previously presented) A method of targeting proteins involved in lipid or fatty acid biosynthesis into liposomes or lipid bodies, which comprises combining the protein-encoding nucleic acids and one of the following sequences to give a joint protein-encoding sequence:
 - 1. a nucleic acid sequence with the sequence shown in SEQ ID NO: 1,
 - nucleic acid sequences which are derived from the nucleic acid sequence shown in SEQ ID NO: 1 as the result of the degeneracy of the generic code,
 - 3. derivatives of the nucleic acid sequence shown in SEQ ID NO: 1 which encode polypeptides with the amino acid sequences shown in SEQ ID NO: 2 and which have at least 60% homology at the amino acid level,
 - 4. a nucleic acid sequence with the sequence shown in SEQ ID NO: 3 or the aminoterminal portion of the coding region of this sequence, and introducing the resulting sequence into a eukaryotic organism.
- 17. (previously presented) A method of targeting proteins involved in lipid or fatty acid biosynthesis into liposomes or lipid bodies, which comprises introducing at least one nucleic acid sequence as claimed in claim 1 or at least one nucleic acid construct comprising said nucleic acid sequence wherein the nucleic acid sequence is linked to one or more regulatory signals into an oil-producing organism.

- 18. (previously presented) A method of producing fatty acids or lipids, which comprises introducing at least one nucleic acid sequence as claimed in claim 1 or at least one nucleic acid construct comprising said nucleic acid sequence wherein the nucleic acid sequence is linked to one or more regulatory signals into an oil-producing organism, growing this organism and isolating the oil contained in the organism.
- 19. (previously presented) A method of producing fatty acids, which comprises introducing at least one nucleic acid sequence as claimed in claim 1 or at least one nucleic acid construct comprising said nucleic acid sequence wherein the nucleic acid sequence is linked to one or more regulatory signals into an oil-producing organism, growing this organism, isolating the oil contained in the organism and liberating the fatty acids.
- 20. (previously presented) A method as claimed in claim 16, wherein the organism is a plant or a eukaryotic microorganism.